

Abdominal aortic aneurysm

Definition

The aorta is the main blood vessel that supplies blood to the abdomen, pelvis, and legs. An abdominal aortic aneurysm occurs when an area of the aorta becomes very large or balloons out.

Alternative Names

Aneurysm - aortic; AAA

Causes

The exact cause of the condition is unknown. Factors that can increase your risk of developing the problem include:

- Smoking
- High blood pressure
- Male gender
- Genetic factors

An abdominal aortic aneurysm is most often seen in males over age 60 who have one or more risk factors. The larger the aneurysm, the more likely it is to break open. This can be life-threatening.

Symptoms

Aneurysms can develop slowly over many years, often with no symptoms. Symptoms may come on quickly if the aneurysm expands rapidly, tears open or leaks blood within the wall of the vessel (aortic dissection).

Symptoms of rupture include:

- Pain in the abdomen or back. The pain may be severe, sudden, persistent, or constant. It may spread to the groin, buttocks, or legs.
- Passing out
- Clammy skin
- Dizziness
- Nausea and vomiting
- Rapid heart rate
- Shock



Exams and Tests

Your doctor will examine your abdomen and feel the pulses in your legs. The doctor may find:

- A lump (mass) in the abdomen
- Pulsating sensation in the abdomen
- Stiff or rigid abdomen

You may have an abdominal aortic aneurysm that is not causing any symptoms. Your doctor may find this problem by doing the following tests:

- Ultrasound of the abdomen when the abdominal aneurysm is first suspected



Abdominal aortic aneurysm

- [CT scan of the abdomen](#) to confirm the size of the aneurysm
- CTA (computed tomographic angiogram) to help with surgical planning

Any one of these tests may be done when you're having symptoms.

Treatment

If you have bleeding inside your body from an aortic aneurysm, you will need [abdominal aortic aneurysm repair](#).

If the aneurysm is small and there are no symptoms:

- Surgery is rarely done.
- You and your doctor must decide if the risk of having surgery is smaller than the risk of bleeding if you do not have surgery.
- Your doctor may want to check the size of the aneurysm with ultrasound tests every 6 months.

Most of the time, surgery is done if the aneurysm is bigger than 2 inches (5.5 cm) across or growing quickly. The goal is to do surgery before complications develop.

There are two types of surgery:

- [Traditional \(open\) repair](#). A large cut is made in your abdomen. The abnormal vessel is replaced with a graft made of man-made material.
- [Endovascular stent grafting](#). This procedure can be done without making a large cut in your abdomen, so you may recover more quickly. This may be a safer approach if you have certain other medical problems. Endovascular repair can sometimes be done for a leaking or bleeding aneurysm.

Outlook (Prognosis)

The outcome is often good if you have surgery to repair the aneurysm before it ruptures.

When an abdominal aortic aneurysm [begins to tear](#) or ruptures, it is a medical emergency. Only about 1 in 5 patients survive a ruptured abdominal aneurysm.

When to Contact a Medical Professional

Go to the emergency room or call 911 if you have pain in your belly or back that is very bad or does not go away.

Prevention

To reduce the risk of aneurysms:

- Eat a heart-healthy diet, exercise, stop smoking (if you smoke), and reduce stress.
- If you have high blood pressure or diabetes, take your medicines as your doctor has told you.

People over age 65 who have ever smoked should have a screening ultrasound done once.

References

Tracci MC, Cherry KJ. The aorta. In: Townsend CM Jr, Beauchamp RD, Evers BM, Mattox KL, eds. *Sabiston Textbook of Surgery*. 19th ed. Philadelphia, Pa: Saunders Elsevier; 2012:chap 62.

Lewiss RE, Egan DJ, Shreves A. Vascular abdominal emergencies. *Emerg Med Clin North Am*. 2011;29(2):253-72, viii.

Greenhalgh RM, Powell JT. Endovascular repair of abdominal aortic aneurysm. *N Engl J Med*. 2008;358:494-501.

[Lederle FA, Kane RL, MacDonald R, Wilt TJ. Systematic review: repair of unruptured abdominal aortic aneurysm. *Ann Intern Med*. 2007;146:735-741.](#)

Braverman AC, Thompson RW, Sanchez LA. Diseases of the aorta. In: Bonow RO, Mann DL, Zipes DP, Libby P, eds. *Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine*. 9th ed. Philadelphia, Pa: Saunders Elsevier; 2011:chap 60.

Review Date: 8/13/2013

Reviewed By: Matthew M. Cooper, MD, MBA, FACS, Medical Director, CareCore National, Bluffton, SC. Review provided by VeriMed Healthcare Network. Also reviewed by David Zieve, MD, MHA, Bethanne Black, and the A.D.A.M. Editorial team.

A.D.A.M., Inc. is accredited by URAC, also known as the American Accreditation HealthCare Commission (www.urac.org). URAC's [accreditation program](#) is an independent audit to verify that A.D.A.M. follows rigorous standards of quality and accountability. A.D.A.M. is among the first to achieve this important distinction for online health information and services. Learn more about A.D.A.M.'s [editorial policy](#), [editorial process](#) and [privacy policy](#). A.D.A.M. is also a founding member of Hi-Ethics and subscribes to the principles of the Health on the Net Foundation (www.hon.ch).