West Kentucky Surgical, Inc Dr. J Craig Dowdy

Colonoscopy Preparation Instructions

Procedure Location: Murray Calloway County Hospital

803 Popular Street Murray KY, 42071.

General Information:

- Should you have any questions regarding your preparation or if you need to reschedule your appointment, please contact our office at 270-753-2444
- Please bring your insurance cards and a list of your medications. Please leave all jewelry, body piercings and valuables at home.

Frequently Asked Questions:

- If I feel nauseated and don't think I can drink more prep, what should I do?
 - It is very important that you drink the entire amount of the solution. You may take a break of 30 minutes and resume. You may also try drinking more slowly and/or use a straw.
- I have taken most of the solution without any results, what should I do?
 - Continue drinking the solution until it's completely gone. It can take up to 2 hours before you start to see results.
- My stools are loose and watery, do I still need to finish the solution?
 - Yes, it is very important to drink all of the solution.
- Can I have gum or hard candy before my procedure?
 - o Yes
- Can I have the colonoscopy if I start my menstrual cycle?
 - o Yes
- Can I take something if I have a headache?
 - Yes, Tylenol or Advil
- Can I use something other than Gatorade to mix the Miralax with?'
 - Yes, as long as it is a CLEAR liquid
- Is it alright if I shower and brush my teeth before my procedure
 - Yes, please shower and brush your teeth prior to your procedure

Preparing For Your Colonoscopy

- Read the instructions in the pamphlet carefully a few days prior to your colonoscopy
- You will need an adult companion to drive you home following your scope
- The Pharmacist can help you with picking out the correct medications if you have questions.

Purchase:

Dulcolax Laxative (Bisacodyl) – 4 tablets Miralax Powder – 238 grams Gatorade – 64oz; No red or purple

The Day Prior to Your Colonoscopy - Please Read Carefully

- 1. You are only able to drink CLEAR liquids. Clear liquids include:
 - a. Water, Gatorade, Tea, Coffee but no creamer, Chicken/Beef broth, Jell-O (no red, fruit, or toppings), any type soda, Cranberry/Apple/White Grape Juice.
- 2. At 12:00 noon the day before your exam, take 4 Dulcolax Laxative (Bisacodyl) tablets with 8 ounces of clear liquid.
- 3. Between 4:00pm and 6:00pm the day prior to your exam, mix the full 238 grams of Miralax powder with the 64oz of Gatorade. Shake the solution until completely dissolved.
- 4. Drink an 8oz glass of the solution every 10-15 minutes until it has all been consumed.
- 5. Drink at least 16 ounces of clear liquid following the prep solution.
- 6. You may continue to drink clear liquids up until 4 hours prior to your scheduled procedure.
- 7. Take your routine medication (including Aspirin and/or Plavix with a sip of water at their regular times.
- 8. Notify our office if you are taking COUMADIN or DIABETIC medication.
- 9. The day prior to your procedure it is important to drink lots of clear liquids.